

Sources

Collins, J. G. (1997). Prevalence of selected chronic conditions: United States 1990–1992. *Vital and Health Statistics*, 10(194). Hyattsville, MD: National Center for Health Statistics

Cruickshanks KJ, Klein R, Klein BEK, Wiley TL, Nondahl DM, Tweed TS. Cigarette Smoking and Hearing Loss: The Epidemiology of Hearing Loss Study. *JAMA*. 1998;279(21):1715–1719.

Hearing Health Foundation. Hearing Loss & Tinnitus Statistics (2016). Retrieved from: <https://hearinghealthfoundation.org/hearing-loss-tinnitus-statistics>

Hull, R. H., & Kerschen, S. R. (2010). The Influence of Cardiovascular Health on Peripheral and Central Auditory Function in Adults. *American Journal*.

National Institute on Deafness and Other Communication Disorders (NIDCD) (2018). Retrieved from: www.nidcd.nih.gov/health/hearing-loss-older-adults

National Institutes of Health (2008). Retrieved from <https://www.nih.gov/news-events/news-releases/hearing-loss-common-people-diabetes>

Patel, A., & Groppo, E. (2010). Management of temporal bone trauma. *Craniofacial trauma & reconstruction*, 3(2), 105–113. <https://doi.org/10.1055/s-0030-1254383>